

संचालनालय स्वास्थ्य सेवायें
मध्य प्रदेश

क्रमांक/अ.प्रशा./रेबीज/2021/638
प्रति,

भोपाल, दिनांक 14/6/2021

समस्त, मुख्य चिकित्सा एवं स्वास्थ्य अधिकारी,
समस्त, सिविल सर्जन सह मुख्य अस्पताल अधीक्षक,
मध्य प्रदेश।

विषय:- कोविड-19 महामारी में एनीमल बाइट एवं रेबीज के केसेस के प्रबंधन के संबंध में।

—00—

विषयांतर्गत कोविड 19 महामारी में एनीमल बाइट एवं रेबीज के केसेस के प्रबंधन के संबंध में राष्ट्रीय रोग नियंत्रक केन्द्र भारत सरकार द्वारा गाईडलाईन्स जारी की गई है, जो पत्र के साथ संलग्न है।

आपको निर्देशित है कि राष्ट्रीय रोग नियंत्रक केन्द्र भारत सरकार द्वारा जारी गाईडलाईन्स अनुसार एनीमल बाइट एवं रेबीज के केसेस का प्रबंधन एवं रेबीज वेक्सीनेशन किया जाना सुनिश्चित करें।

संलग्न :- गाईडलाईन्स।

11/06/2021
(डॉ.सतोष जैन)

अपर संचालक, (अ.प्र.)
संचाल नालय स्वास्थ्य सेवायें
मध्य प्रदेश

पृ.क्रमांक./अ.प्रशा./ रेबीज /2021/635
प्रतिलिपि :- सूचनार्थ एवं आवश्यक कार्यवाही हेतु ।

भोपाल ,दिनांक 14/6/2021

01. अपर मुख्य सचिव, लोक स्वास्थ्य एवं परिवार कल्याण विभाग वल्लभ भवन मंत्रालय।
02. आयुक्त स्वास्थ्य स्थानीय कार्यालय।
03. मिशन संचालक, एन.एच.एम. भोपाल मध्य प्रदेश।
04. समस्त क्षेत्रीय संचालक, स्वास्थ्य सेवायें मध्य प्रदेश ।

11/06/2021
अपर संचालक, (अ.प्र.)
संचाल नालय स्वास्थ्य सेवायें
मध्य प्रदेश



भारत सरकार
राष्ट्रीय रोग नियंत्रक केन्द्र
स्वास्थ्य सेवा महानिदेशालय
22, शाम नाथ मार्ग, दिल्ली-110054
Government of India

NATIONAL CENTER FOR DISEASE CONTROL
[Formally Known as National Institute of Communicable Disease (NICD)]

Directorate General of Health Services
Ministry of Health & Family Welfare, Gov. of India
22, Sham Nath Marg, Delhi-110054



Direct: 00-91-11-23913148
23922132, 23944035
Fax: 00-91-11-23922677
Email: dirnicd@nic.in
Website: www.ncdc.gov.in
www.idsp.nic.in

File No: NRCP/55/92/05/2021/DZDP-NCDC

Date 10 June 2021

Dear All,

As it is understood that in view of COVID- 19 pandemic, a range of complex issues have evolved with regard to diagnosis, treatment and vaccination for different diseases including Rabies.

As per pervious communication from the Ministry vide DO: T-22020/14/2020 Dated- 14/1/2021 it is advised that if co-administration of COVID- 19 vaccine is required with other vaccine, then there should be a gap of at least 14 days between two vaccines. However COVID- 19 and other vaccines may be administered within a shorter period in situation where the benefits of vaccination are deemed to outweigh the potential unknown risks of vaccine co-administration.

Hence in view of current COVID- 19 pandemic following is advised for management of animal bites and rabies cases

1. All animal bite cases should continue to receive the Rabies vaccination as per Rabies Post -Exposure Prophylaxis schedule as per National Guidelines irrespective of COVID- 19 status of Individual (as Rabies is 100 % percent fatal disease and its benefits outweigh the risk associated with it). However, clinicians are advised to use their own discretion (with risk verses benefits) in individual cases of Immuno- compromised cases, patients with serious life-threatening conditions etc.
2. If COVID-19 vaccines are administered within 14 days of Rabies vaccine, doses need not to be repeated and schedule of vaccination for either vaccine should be completed as per National Guidelines.
3. All Rabies cases should also receive due priority for admission in respective Infectious disease hospital/Medical colleges and standard care, as appropriate should be administered to all such cases.

For more information on rabies vaccines and animal bite management you are advised to refer to National Guidelines of Rabies prophylaxis and management (2019) at <https://www.ncdc.gov.in/index1.php?lang=1&level=1&sublinkid=146&lid=150>

This issues with approval DGHS, MoH&FW, GOI & meant for necessary circulation with respective nodal officers of all Anti-Rabies clinics and hospital please.

With Regards

Dr. Simmi Tiwari
Joint Director & In charge of
National Rabies Control Programme

To Nodal officers under NRCP- All States/UTs

Copy to

- PS to Director NCDC
- PS to DGHS, MOHFW, Govt of India



Antibiotic Resistance Containment Stewardship: Our Role, Our Responsibility
Judicious Use of Antibiotic: Key to Contain Antibiotic Resistance

